## **Staying Healthy: Recommendations for WOMEN**



The Veterans Health Administration (VHA) aims to help you stay healthy. The table below lists the preventive health services (screening tests, medications, health counseling, and vaccines) that VHA recommends. Screening tests are used to look for health conditions before there are symptoms. These recommendations apply only to adult women of average risk. You are a woman of average risk if you have no personal or family history or symptoms of the conditions listed below. If you are having symptoms of a condition, please talk with your provider.

Which preventive health services are recommended for average risk women in my age group?

Green: Recommended Yellow: Recommended for some women – talk with your provider Red: Not recommended (NR) SCREENING TESTS FOR AVERAGE RISK VETERANS **HEALTH CONDITIONS** 70–79 years 80 years and older 18-29 years 40-49 years 50-59 years 60-69 years 30–39 years Talk with your Recommended every two years (age 40-74) **Breast Cancer** Talk with your provider Not recommended provider Requests for annual screening can be honored after discussion of risks and benefits (age 75 and older) (age 18–29) \*Service Act Veterans (age 30-39)\* Every 3 years Recommended. Not recommended for those with prior adequate screening **Cervical Cancer** (age 21–29) Every 3 or 5 years, depending on tests chosen (age 30–65) (age 66 and older) Talk with your provider NR (age 86 Recommended. Not recommended (age 18-44) **Colon Cancer** Frequency varies by test chosen (age 45–75) (age 76–85) and older) **Depression** Recommended every year Recommended for some women — talk with your provider (age 18 and older) **Hepatitis B Infection** Recommended in adults aged 18-79 years, and in all pregnant women, during each pregnancy **Hepatitis C Infection High Blood Pressure** Recommended every year (age 18 and older) Recommended once (age 18 and older); in all pregnant Veterans; annually in Veterans with ongoing risk factors **HIV Infection** Recommended for some women — talk with your Not recommended Not recommended (age 18–49) **Lung Cancer** (age 81 and older) provider (age 50–80) Recommended for some women — talk with your provider (age 18—64) Recommended once (age 65 and older) **Osteoporosis** Recommended. Test for gonorrhea and Recommended for some women, talk with your provider about testing for gonorrhea, chlamydia, and syphilis (age 25 and older) chlamydia (age 18-24). **Sexually Transmitted Infections** Talk with your Recommended for all pregnant Veterans, early screening for syphilis infection, regardless of age. provider about syphilis testing. **MEDICATIONS** 18-29 years 80 years and older 30-39 years 40-49 years 50-59 years 60-69 years 70-79 years **HEALTH CONDITIONS** Recommended. Daily folic acid supplement for any woman who may become **Folic Acid for Pregnancy Planning** Not recommended after child-bearing age (age 55 and older) pregnant (age 18-55) **Aspirin to Prevent** Not recommended (age 18–39) Not recommended (age 60 and older) Talk with your provider (age 40–59) **Cardiovascular Disease** Talk with your provider (age 19 and older) **PrEP to Prevent HIV Infection Statin to Prevent** Not recommended (age 19–40) Recommended for some women — talk with your provider (age 40 and older) **Cardiovascular Disease HEALTH COUNSELING HEALTH CONDITIONS** 18-29 years 30-39 years 40-49 years 50-59 years 70-79 years 80 years and older 60–69 years Recommended every visit (if using tobacco) (age 18 and older) **Tobacco Use** Talk with your provider about healthy alcohol use (age 18 and older) **Alcohol Use** Talk with your provider about a healthy diet and physical activity (age 18 and older) **Healthy Diet and Physical Activity Overweight & Obesity** Recommended every year (age 18 and older) **VACCINES HEALTH CONDITIONS** 80 years and older 19-29 years 30-39 years 40-49 years 50-59 years 60-69 years 70–79 years Recommended for all women — talk with your provider to make sure you stay up-to-date (<a href="https://www.va.gov/health-care/covid-19-vaccine/">https://www.va.gov/health-care/covid-19-vaccine/</a>) **COVID-19** Recommended for some women — talk with your provider (age 19 and older) **Hepatitis A Hepatitis B** Recommended for all women who have not already had Hepatitis B vaccine (age 19-59) Recommended for some women - talk with your provider (age 60 and older) **Herpes Zoster (Shingles)** Recommended for some women — talk with your provider (age 19 and older) Recommended 2 doses (age 50 and older), with second dose 2–6 months after first dose RZV vaccine 2–3 doses (age **Human Papilloma Virus (HPV)** Not recommended (age 46 and older) 19–26) if series not Talk with your provider (age 27–45) HPV9 vaccine completed Recommended every year (age 19 and older) Influenza (Flu) Injectable, Inhaled Measles, Mumps, and Recommended for some women — talk with your provider (age 19 and older) Rubella (MMR) Recommended for some women — talk with your provider (age 19 and older) **Meningococcal Disease Pneumococcal Disease** Recommended — talk with your provider to make sure PCV20 vaccine, PCV15 vaccine, Recommended for some women — talk with your provider (age 19—64) you are up-to-date (age 65 and older) PPSV23 vaccine "One time" (single lifetime) dose "One time" (single lifetime) dose **Respiratory Syncytial Virus (RSV) Pregnant Veterans** are recommended to receive a one-time (single lifetime) RSV Abrysvo® recommended for women age 75 recommended for some women – vaccine between 32-36 weeks gestation during RSV season RSV vaccine talk with your provider (age 60–74) and older Recommend at least 1 dose. Get additional dose with every pregnancy **Tdap (Tetanus, Diphtheria** (Or for wound management if > 5 years have passed since last tetanus toxoid-containing booster vaccine). & Pertussis) Recommended. Booster every 10 years (age 19 and older) (Either Tdap or Td may be used) **Td (Tetanus and diphtheria)** (Or for wound management if > 5 years have passed since last tetanus toxoid-containing booster vaccine). Recommended for some women - talk with your Varicella (Chickenpox) provider (2 doses if born 1980 or later)

Current evidence is based on birth sex. Because there is limited research on transgender and gender-diverse people, VA recommends talking with your health care provider for individualized clinical preventive services to achieve optimal health and well-being. See <a href="https://www.patientcare.va.gov/LGBT/">https://www.patientcare.va.gov/LGBT/</a>





