

Staying Healthy: Recommendations for WOMEN



U.S. Department of Veterans Affairs Veterans Health Administration

The Veterans Health Administration (VHA) aims to help you stay healthy. The table below lists the **preventive health services** (screening tests, medications, health counseling, and vaccines) that VHA recommends. Screening tests are used to look for health conditions before there are symptoms. These recommendations apply only to adult women of **average risk**. You are a woman of average risk if you have **no personal or family history or symptoms** of the conditions listed below. If you are having symptoms of a condition, please talk with your provider.

Which preventive health services are recommended for average risk women in my age group?

Green: Recommended Yellow: Recommended for some men – talk with your provider

Red: Not recommended (NR)

SCREENING TESTS FOR AVERAGE RISK VETERANS

HEALTH CONDITIONS	18–29 years	30–39 years	40–49 years	50–59 years	60–69 years	70–79 years	80 years and older			
Breast Cancer		mmended 18—39)	Recommend biennia Have op	Recommended annually (45-54) (55 and older or have opportunity to continue annually) ortunity to begin annual screening (40-44) Recommended for some women – talk with your provider (age 75 and older)						
Cervical Cancer	Every 3 years (age 21-29)Recommended.Not recommended (age 66 ar with prior adequateEvery 3 or 5 years, depending on tests chosen (age 30-65)with prior adequate									
Colon Cancer	Not recor	nmended (age 18—44)		Recomm Frequency varies by te			Talk with your provider (age 76–85)NR (age 86 and older)			
Depression	Recommended every year									
Hepatitis B Infection	Recommended for some women — talk with your provider (age 18 and older)									
Hepatitis C Infection	Recommended in adults aged 18-79 years, and in all pregnant women, during each pregnancy									
High Blood Pressure	Recommended every year (age 18 and older)									
HIV Infection	Recommended once (age 18 and older); in all pregnant Veterans; annually in Veterans with ongoing risk factors									
Lung Cancer	Not recommended (age 18–49) Recommended for some women – talk with your provider (age 50–80) Not recommended (age 81 and older)									
Osteoporosis	Recommended for some women – talk with your provider (age 18–64) Recomm					commended once (ag	je 65 and older)			
Sexually Transmitted	nitted Recommended. Test for gonorrhea and chlamydia (age 18–24). Recommended for some women, talk with your provider about testing for gonorrhea, chlamydia, and syphilis (age 25 a net testing for gonorrhea, chlamydia, and syphilis (age 25 a									
Infections	Talk with your provider about syphilis testing.	Recor	nmended for all preg	nant Veterans, early scre	eening for syphilis infec	ction, regardless of a	je.			

MEDICATIONS

HEALTH CONDITIONS	18–29 years	30–39 years	40–49 years	50-59	years	60–69 years	70–79 years	80 years and older
Folic Acid for Pregnancy Planning	Recommended. Daily folic acid supplement for any woman who may become pregnant (age 18-55) Not recommended after child-bearing age (age 55 and older)							e 55 and older)
Aspirin to Prevent Cardiovascular Disease	Not recommended (age 18–39) Talk with your provider (age 40–59) Not recommended (age 60 and older)						nd older)	
PrEP to Prevent HIV Infection	Talk with your provider (age 19 and older)							
Statin to Prevent Cardiovascular Disease	Not recommended (age 19–40) Recommended for some women – talk with your provider (age 40 and older)							

HEALTH COUNSELING

HEALTH CONDITIONS	18–29 years	30–39 years	40–49 years	50–59 years	60–69 years	70–79 years	80 years and older			
Tobacco Use	Recommended every visit (if using tobacco) (age 18 and older)									
Alcohol Use	Talk with your provider about healthy alcohol use (age 18 and older)									
Healthy Diet and Physical Activity	Talk with your provider about a healthy diet and physical activity (age 18 and older)									
Overweight & Obesity	Recommended every year (age 18 and older)									

VACCINES

HEALTH CONDITIONS	19–29 years	30–39 years	40–49 years	50–59 years	60–69 years	70–79 years	80 years and older			
COVID-19	Recommended for all women - talk with your provider to make sure you stay up-to-date (https://www.va.gov/health-care/covid-19-vaccine/)									
Hepatitis A	Recommended for some women – talk with your provider (age 19 and older)									
Hepatitis B	Recommended for	Recommended for all women who have not already had Hepatitis B vaccine (age 19-59) 60 and older)								
Herpes Zoster (Shingles) RZV vaccine	Recommended	for some women – talk (age 19 and older)	with your provider	Reco	mmended 2 doses (age 50 and older), with second dose 2–6 months after first dose					
Human Papilloma Virus (HPV) HPV9 vaccine	2–3 doses (age 19–26) if series Talk with your provider (age 27–45) Not recommended (age 46 and older) not completed									
Influenza (Flu) Injectable, Inhaled		Recommended every year (age 19 and older)								
Measles, Mumps, and Rubella (MMR)	Recommended for some women – talk with your provider (age 19 and older)									
Meningococcal Disease	Recommended for some women – talk with your provider (age 19 and older)									
Pneumococcal Disease PCV20 vaccine, PCV15 vaccine, PPSV23 vaccine	Recommended for some women – talk with your provider (age 19–64) Recommended - talk with your provider to make sure you are up-to-date (age 65 and older)									
Respiratory Syncytial Virus (RSV) RSV vaccine	Pregnant Veterans are recommended to receive a one-time (single lifetime) RSV Abrysvo® vaccine between 32-36 weeks gestation during RSV season during recommended for some women – talk with your provider (age 60–74) "One time" (single lifetime)						e recommended for			
Tdap (Tetanus, Diphtheria & Pertussis)	Recommend at least 1 dose. Get additional dose with every pregnancy (Or for wound management if > 5 years have passed since last tetanus toxoid-containing booster vaccine).									
Td (Tetanus and diphtheria)	Recommended. Booster every 10 years (age 19 and older) (Either Tdap or Td may be used) (Or for wound management if > 5 years have passed since last tetanus toxoid-containing booster vaccine).									
Varicella (Chickenpox)		for some women - talk w 2 doses if born 1980 or lat								

Current evidence is based on birth sex. Because there is limited research on transgender and gender-diverse people, VA recommends talking with your health care provider for individualized clinical preventive services to achieve optimal health and well-being. See <u>https://www.patientcare.va.gov/LGBT/</u>



To view a copy on your mobile device, scan the QR code or visit:

https://www.prevention.va.gov/Healthy_Living/Get_Recommended_Screening_Tests_and_Immunizations_for_Women.asp Talk with your health care team during your next appointment to ensure you are up to date on your preventive care.