

# Staying Healthy: Recommendations for WOMEN



The Veterans Health Administration (VHA) aims to help you stay healthy. The table below lists the **preventive health services** (screening tests, medications, health counseling, and vaccines) that VHA recommends. Screening tests are used to look for health conditions before there are symptoms. These recommendations apply only to adult women of **average risk**. You are a woman of average risk if you have **no personal or family history or symptoms** of the conditions listed below. If you are having symptoms of a condition, please talk with your provider.

# Which preventive health services are recommended for average risk women in my age group?

Green: Recommended Yellow: Recommended for some men – talk with your provider Red: Not recommended (NR)

#### **SCREENING TESTS FOR AVERAGE RISK VETERANS**

| <b>HEALTH CONDITIONS</b>           | 18–29 years  | 30–39 years  | 40-49 years | 50–59 years | 60–69 years | 70–79 years   | 80 years and older |  |  |  |  |
|------------------------------------|--|--|-------------|-------------|-------------|---|--------------------|--|--|--|--|
| Breast Cancer                      |  | recommended Recommended annually (45-54 Recommend biennial (55 and older or have opportunit Have opportunity to begin annual screeni |             |             |             | y to continue annually) — talk with your provider         |                    |  |  |  |  |
| Cervical Cancer                    | Every 3 years<br>(age 21–29)   |  |             |             |             |   |                    |  |  |  |  |
| Colon Cancer                       | Not rec  | Not recommended (age 18–44)  Recommende Frequency varies by test cho   |             |             |             | Talk with your NR (age 86 provider (age 76–85) and older) |                    |  |  |  |  |
| Depression                         | Recommended every year   |  |             |             |             |   |                    |  |  |  |  |
| Hepatitis B Infection              | Recommended for some women — talk with your provider (age 18 and older)  |  |             |             |             |   |                    |  |  |  |  |
| Hepatitis C Infection              | Recommended in adults aged 18-79 years, and in all pregnant women, during each pregnancy                                       |  |             |             |             |   |                    |  |  |  |  |
| High Blood Pressure                | Recommended every year (age 18 and older)  |  |             |             |             |   |                    |  |  |  |  |
| HIV Infection                      | Recommended once (age 18 and older); in all pregnant Veterans; annually in Veterans with ongoing risk factors                  |  |             |             |             |   |                    |  |  |  |  |
| Lung Cancer                        | Not recommended (age 18–49)  Recommended for some women — talk with your provider (age 50–80)  Not recommended (age 81 and old |  |             |             |             |   |                    |  |  |  |  |
| Osteoporosis                       | Recommended for some women — talk with your provider (age 18—64)  Recommended once   |  |             |             |             |   | ige 65 and older)  |  |  |  |  |
| Sexually Transmitted<br>Infections | Recommended.<br>Test for<br>gonorrhea and<br>chlamydia (age<br>18–24).   | Recommended for some women, talk with your provider about testing for gonorrhea, chlamydia, and syphilis (age 25 and older)          |             |             |             |   |                    |  |  |  |  |
|                                    | Talk with<br>your provider<br>about syphilis<br>testing.   | Recommended for all pregnant Veterans, early screening for syphilis infection, regardless of age.                                    |             |             |             |   |                    |  |  |  |  |

# **MEDICATIONS**

| HEALTH CONDITIONS                               | 18–29 years  | 30–39 years | 40–49 years | 50-59 | .59 years 60–69 years |                    | 70-79 years | 80 years and older |
|---|--|-------------|-------------|-------|-----------------------|--------------------|-------------|--------------------|
| Folic Acid for Pregnancy<br>Planning            | Recommended. Daily folic acid supplement for any woman who may become pregnant (age 18-55)  Not recommended after child-bearing age (age 55 and older) |             |             |       |                       |                    |             |                    |
| Aspirin to Prevent<br>Cardiovascular<br>Disease | Not recommended (age 18–39)  Talk with your provider (age 40–59)  Not recommended (  |             |             |       |                       | ommended (age 60 a | nd older)   |                    |
| PrEP to Prevent<br>HIV Infection                | Talk with your provider (age 19 and older)   |             |             |       |                       |                    |             |                    |
| Statin to Prevent<br>Cardiovascular Disease     | Not recommended (age 19–40)  Recommended for some women – talk with your provider (age 40 and older)   |             |             |       |                       |                    |             |                    |

## **HEALTH COUNSELING**

| HEALTH CONDITIONS                     | 18–29 years   | 30–39 years | 40-49 years | 50-59 years | 60–69 years | 70–79 years | 80 years and older |  |  |  |  |
|---------------------------------------|---|-------------|-------------|-------------|-------------|-------------|--------------------|--|--|--|--|
| Tobacco Use                           | Recommended every visit (if using tobacco) (age 18 and older)                         |             |             |             |             |             |                    |  |  |  |  |
| Alcohol Use                           | Talk with your provider about healthy alcohol use (age 18 and older)                  |             |             |             |             |             |                    |  |  |  |  |
| Healthy Diet and<br>Physical Activity | Talk with your provider about a healthy diet and physical activity (age 18 and older) |             |             |             |             |             |                    |  |  |  |  |
| Overweight & Obesity                  | Recommended every year (age 18 and older)   |             |             |             |             |             |                    |  |  |  |  |

## **VACCINES**

|   |   |   |             |             |   |         |      | 1  |  |  |
|---|---|---|-------------|-------------|---|---------|------|--|--|--|
| HEALTH CONDITIONS   | 19–29 years   | 30–39 years   | 40–49 years | 50-59 years | 60–69 years   | 70-79 y | ears | 80 years and older   |  |  |
| COVID-19  | Recommended for all women — talk with your provider to make sure you stay up-to-date (https://www.va.gov/health-care/covid-19-vaccine/)   |   |             |             |   |         |      |  |  |  |
| Hepatitis A   | Recommended for some women — talk with your provider (age 19 and older)   |   |             |             |   |         |      |  |  |  |
| Hepatitis B   | Recommended for all women who have not already had Hepatitis B vaccine (age 19-59)  Recommended for some women - talk with your provide 60 and older)   |   |             |             |   |         |      | ith your provider (age   |  |  |
| Herpes Zoster<br>(Shingles) RZV vaccine                                 | Recommended for some women — talk with your provider (age 19 and older)   |   |             |             | Recommended 2 doses (age 50 and older), with second dose 2–6 months after first dose                        |         |      |  |  |  |
| Human Papilloma<br>Virus (HPV)<br>HPV9 vaccine                          | 2–3 doses (age<br>19–26) if series<br>not completed  Talk with your provider (age 27–45)  Not recommended (age 46 and older)  |   |             |             |   |         |      |  |  |  |
| Influenza (Flu)<br>Injectable, Inhaled                                  | Recommended every year (age 19 and older)   |   |             |             |   |         |      |  |  |  |
| Measles, Mumps, and<br>Rubella (MMR)                                    | Recommended for some women — talk with your provider (age 19 and older)   |   |             |             |   |         |      |  |  |  |
| Meningococcal Disease   | Recommended for some women — talk with your provider (age 19 and older)   |   |             |             |   |         |      |  |  |  |
| Pneumococcal Disease<br>PCV20 vaccine, PCV15<br>vaccine, PPSV23 vaccine | Recommended for some women — talk with your provider (age 19—64)  Recommended - talk with your provider to make sure you are up-to-date (age 65 and older)  |   |             |             |   |         |      |  |  |  |
| Respiratory Syncytial<br>Virus (RSV)<br>RSV vaccine                     | <b>Pregnant Veterans</b> are recommended to receive a one-time (single lifetim RSV Abrysvo® vaccine between 32-36 weeks gestation during RSV season   |   |             |             | "One time" (single lifetime)<br>dose recommended for<br>some women – talk with<br>your provider (age 60–74) |         | dose | ime" (single lifetime)<br>recommended for<br>en age 75 and older |  |  |
| Tdap (Tetanus,<br>Diphtheria & Pertussis)                               | Recommend at least 1 dose. Get additional dose with every pregnancy (Or for wound management if $>$ 5 years have passed since last tetanus toxoid-containing booster vaccine).                    |   |             |             |   |         |      |  |  |  |
| Td (Tetanus and diphtheria)   | Recommended. Booster every 10 years (age 19 and older) (Either Tdap or Td may be used) (Or for wound management if $>$ 5 years have passed since last tetanus toxoid-containing booster vaccine). |   |             |             |   |         |      |  |  |  |
| Varicella (Chickenpox)  |   | d for some women - talk w<br>(2 doses if born 1980 or lat |             |             |   |         |      |  |  |  |

Current evidence is based on birth sex. Because there is limited research on transgender and gender-diverse people, VA recommends talking with your health care provider for individualized clinical preventive services to achieve optimal health and well-being. See <a href="https://www.patientcare.va.gov/LGBT/">https://www.patientcare.va.gov/LGBT/</a>



To view a copy on your mobile device, scan the QR code or visit:

https://www.prevention.va.gov/Healthy\_Living/Get\_Recommended\_Screening\_Tests\_and\_Immunizations\_for\_Women.asp Talk with your health care team during your next appointment to ensure you are up to date on your preventive care.