Staying Healthy: **Recommendations for WOMEN**



The Veterans Health Administration (VHA) aims to help you stay healthy. The table below lists the preventive health services (screening tests, medications, health counseling, and vaccines) that VHA recommends. Screening tests are used to look for health conditions before there are symptoms. These recommendations apply only to adult women of average risk. You are a woman of average risk if you have no personal or family history or symptoms of the conditions listed below. If you are having symptoms of a condition, please talk with your provider.

Which preventive health services are recommended for average risk women in my age group?

Green:	Recommended
Yellow:	Recommended for some women – talk with your provider
Red:	Not recommended (NR)

REENING TESTS FOR AVERAGE RISK VETERANS

HEALTH CONDITIONS	18–29 years	30–39 years	40–49 years	50–59 years	60–69 years	70–79 years	80 years a	and older		
Breast Cancer		nmended 8–39)	Recommended annually (45-54) Recommend biennial (55 and older or have opportunity to continue annually) Have opportunity to begin annual screening (40-44)				Recommended for some women — talk with your provider (age 75 and older)			
Cervical Cancer	Every 3 years (age 21–29)	Every 3 o		Recommended. pending on tests chosen (age 30–65) Not recommended.			ded for those with prior adequate screening (age 66 and older)			
Colon Cancer	Not reco	mmended (age 18–44)		Recommended. Frequency varies by test chosen (age 45–75)			Talk with your provider NR (age 86 and older)			
Depression		Recommended every year								
Hepatitis B Infection		Recommended for some women — talk with your provider (age 18 and older)								
Hepatitis C Infection	Recommended in adults aged 18-79 years, and in all pregnant women, during each pregnancy									
High Blood Pressure	Recommended every year (age 18 and older)									
HIV Infection	Recommended once (age 18 and older); in all pregnant Veterans; annually in Veterans with ongoing risk factors									
Lung Cancer	Not recommended (age 18 //U)					Not recom (age 81 ar				
Osteoporosis	Recommended for some women — talk with your provider (age 18—64) Recommended for some women — talk with your provider (age 18—64) Recommended for some women — talk with your provider (age 18—64)					nge 65 and older)				
Sexually Transmitted Infections	Recommended. Test for gonorrhea and chlamydia (age 18–24). Recommended for some women, talk with your provider about testing for gonorrhea, chlamydia, and syphilis (age 25 and older)									
	Talk with your provider about Recommended for all pregnant Veterans, early screening for syphilis infection, regardless of age. syphilis testing.									

MEDICATIONS

HEALTH CONDITIONS	18–29 years	30–39 years	40–49 years	50–59 years	60–69 years	70-79 years	80 years and older
Folic Acid for Pregnancy Planning	Recommended. Daily folic acid supplement for any woman who may become pregnant (age 18-55)			Not recommended after child-bearing age (age 55 and older)			
Aspirin to Prevent Cardiovascular Disease	Not recommended (age 18–39) Talk with your provider (age 40–59)			Not recommended (age 60 and older)			
PrEP to Prevent HIV Infection	Talk with your provider (age 19 and older)						
Statin to Prevent Cardiovascular Disease	Not recommended (age 19–40) Recommended for some women — talk with your provider (age 40 and older)						

HEALTH COUNSELING

HEALTH CONDITIONS	18–29 years	30–39 years	40–49 years	50-59 years	60–69 years	70–79 years	80 years and older
Tobacco Use			Recommended (every visit (if using tobacco)	(age 18 and older)		
Alcohol Use			Talk with your prov	ider about healthy alcohol u	ise (age 18 and older)		
Healthy Diet and Physical Activity			Talk with your provider abo	out a healthy diet and physic	cal activity (age 18 and older	r)	
Overweight & Obesity			Recomi	mended every year (age 18 a	and older)		

Varicella (Chickenpox)

VACCINES												
HEALTH CONDITIONS	19–29 years	30–39 years	40–49 years	50–59 years	60–69 years	70–79 years	80 years and olde					
COVID-19		Recommended for all women — talk with your provider to make sure you stay up-to-date (https://www.va.gov/health-care/covid-19-vaccine/)										
Hepatitis A		Recommended for some women — talk with your provider (age 19 and older)										
Hepatitis B	Recommende	Recommended for all women who have not already had Hepatitis B vaccine (age 19-59) Recommended for some women - talk with your provider (age 60 and older)										
Herpes Zoster (Shingles) RZV vaccine	Recommended for sor	Recommended for some women — talk with your provider (age 19 and older) Recommended 2 doses (age 50 and older), with second dose 2—6 months after first dose										
Human Papilloma Virus (HPV) HPV9 vaccine	2–3 doses (age 19–26) if series not completed	Talk with your provider (a	ge 27–45)	Not recommended (age 46 and older)								
Influenza (Flu) Injectable, Inhaled	Recommended every year (age 19 and older)											
Measles, Mumps, and Rubella (MMR)	Recommended for some women — talk with your provider (age 19 and older)											
Meningococcal Disease			Recommended for some	women — talk with your p	orovider (age 19 and older)							
Pneumococcal Disease PCV20 vaccine, PCV15 vaccine, PPSV23 vaccine	Recommended for some women — talk with your provider (age 19—64) you are up-to-date (age 65 and older)						•					
Respiratory Syncytial Virus (RSV) RSV vaccine	Pregnant Veterans are recommended to receive RSV Abrysvo® vaccine between 32-36 weeks gestation during RSV season Recommended for some women — talk with your provider (age 60 and ol											
Tdap (Tetanus, Diphtheria & Pertussis)		Recommend at least 1 dose. Get additional dose with every pregnancy (Or for wound management if $>$ 5 years have passed since last tetanus toxoid-containing booster vaccine).										
Td (Tetanus and diphtheria)		Recommended. Booster every 10 years (age 19 and older) (Either Tdap or Td may be used) (Or for wound management if $>$ 5 years have passed since last tetanus toxoid-containing booster vaccine).										
Varicella (Chickenpox)	Recommended	d for some women - talk with	your									



provider (2 doses if born 1980 or later)