

Staying Healthy: Recommendations for MEN



The Veterans Health Administration (VHA) aims to help you stay healthy. The table below lists the **preventive health services** (screening tests, medications, health counseling, and vaccines) that VHA recommends. Screening tests are used to look for health conditions before there are symptoms. These recommendations apply only to adult men of **average risk**. You are a man of average risk if you have **no personal or family history or symptoms** of the conditions listed below. If you are having symptoms of a condition, please talk with your provider.

Which preventive health services are recommended for average risk men in my age group?

Green: Recommended

Yellow: Recommended for some men – talk with your provider

Red: Not recommended (NR)

SCREENING TESTS FOR AVERAGE RISK VETERANS

HEALTH CONDITIONS	18–29 years	30–39 years	40–49 years	50-59 years	60–69 years	70–79 y	years 80 years and		
Abdominal Aortic Aneurysm (AAA)	Not recommended (and 18-6/1)					Not recomm (age 76 and			
Colon Cancer	Not recom	mended (age 18–44)		Recommended. Frequency varies by test chosen (age 45–75) Talk with your provider (age 76–85) and					
Depression	Recommended every year								
Hepatitis B Infection	Recommended for some men — talk with your provider (age 18 and older)								
Hepatitis C Infection	Recommended in adults aged 18-79 years								
High Blood Pressure	Recommended every year (age 18 and older)								
HIV Infection	Recommended once (age 18 and older); annually in Veterans with ongoing risk factors								
Lung Cancer	Not recommended (age 18–49)					commended 1 and older)			
Prostate Cancer		Not recommended (a	ge 18–54)		vith your provider (age 55—69)	Not recom	nmended (age 70	and older)	
Sexually Transmitted Infections	Talk with your provider about testing for syphilis and other sexually transmitted infections (age 18 and older)								

MEDICATIONS

HEALTH CONDITIONS	18–29 years	30–39 years	40–49 years	50–59 years	60–69 years	70-79 years	80 years and older	
Aspirin to Prevent Cardiovascular Disease	Not recommend	ded (age 18–39)	Talk with your pro	vider (age 40–59)	Not recommended (age 60 and older)			
PrEP to Prevent HIV Infection	Talk with your provider (age 19 and older)							
Statin to Prevent Cardiovascular Disease	Not recommended (age 19–40) Recommended for some men – talk with your provider (age 40 and older)					lder)		

HEALTH COUNSELING

HEALTH CONDITIONS	18–29 years	30–39 years	40-49 years	50-59 years	60–69 years	70-79 years	80 years and older		
Tobacco Use	Recommended every visit (if using tobacco) (age 18 and older)								
Alcohol Use	Talk with your provider about healthy alcohol use (age 18 and older)								
Healthy Diet and Physical Activity	Talk with your provider about a healthy diet and physical activity (age 18 and older)								
Overweight & Obesity	Recommended every year (age 18 and older)								

VACCINES

HEALTH CONDITIONS	19–29 years	30–39 years	40–49 year	s 50–59 years	60–69 years	70-79	9 years	80 years and older	
COVID-19	Recommended for all men — talk with your provider to make sure you stay up-to-date (https://www.va.gov/health-care/covid-19-vaccine/)								
Hepatitis A	Recommended for some men — talk with your provider (age 19 and older)								
Hepatitis B	Recommended for all men who have not already had Hepatitis B vaccine (age 19-59) Recommended for some men - talk with your provider (age 60 and older)							vith your provider	
Herpes Zoster (Shingles) RZV vaccine	Recommend	ded for some men – talk w (age 19 and older)	ith your provider	Reco	ommended 2 doses (age 50 and older), with second dose 2–6 months after first dose				
Human Papilloma Virus (HPV) HPV9 vaccine	2–3 doses (age 19–26) if series not completed Not recommended (age 46 and older)								
Influenza (Flu) Injectable, Inhaled	Recommended every year (age 19 and older)								
Measles, Mumps, and Rubella (MMR)	Recommended for some men — talk with your provider (age 19 and older)								
Meningococcal Disease	Recommended for some men — talk with your provider (age 19 and older)								
Pneumococcal Disease PCV20 vaccine, PCV15 vaccine, PPSV23 vaccine	Recommended for some men — talk with your provider (age 19—64) Recommended — talk with your provider to make sure you are up-to-date (age 65 and older)								
Respiratory Syncytial Virus (RSV) RSV vaccine	Not recommended (age 18–59)				Recommende men – talk provider (a	with your		mmended for men ge 75 and older	
Tdap (Tetanus, Diphtheria & Pertussis)	Recommend at least 1 dose (age 19 and older) Booster every 10 years (age 19 and older)(Either Tdap or Td may be used) (Or for wound management if > 5 years have passed since last tetanus toxoid-containing booster vaccine).								
Td (Tetanus and diphtheria)	Recommended. Booster every 10 years (age 19 and older) (Either Tdap or Td may be used) (Or for wound management if $>$ 5 years have passed since last tetanus toxoid-containing booster vaccine).								
Varicella (Chickenpox)		ed for some men - talk wit (2 doses if born 1980 or lat							