# Staying Healthy: Recommendations for WOMEN



The Veterans Health Administration (VHA) aims to help you stay healthy. The table below lists the **preventive health services** (screening tests, medications, health counseling, and vaccines) that VHA recommends. Screening tests are used to look for health conditions before there are symptoms. These recommendations apply only to adult women of **average risk**. You are a woman of average risk if you have **no personal or family history or symptoms** of the conditions listed below. If you are having symptoms of a condition, please talk with your provider.

## Which preventive health services are recommended for average risk women in my age group?

Green: Recommended

Yellow: Recommended for some women – talk with your provider

Red: Not recommended (NR)

### **SCREENING TESTS FOR AVERAGE RISK VETERANS**

HEALTH CONDITIONS	18–29 years	30–39 years	40–49 years	50-59 years	60–69 years	<b>70–79</b> y	ears 80 years and older			
Breast Cancer		nmended 18–39)	Recommended annually (45-54) Recommend biennial (55 and older or have opportunity to continue annually) Have opportunity to begin annual screening (40-44)			ıally)	Recommended for some women — talk with your provider (age 75 and older)			
Cervical Cancer	Every 3 years (age 21–29)	Every 3 c	Recommended or 5 years, depending on tes	Not red	Not recommended for those with prior adequate screening (age 66 and older)					
Colon Cancer	Not reco	mmended (age 18–44)	Recommended. Frequency varies by test chosen (age 45–75)				Talk with your prov (age 76–85)	ider NR (age 86 and older)		
Depression		Recommended every year								
Hepatitis B Infection		Recommended for some women — talk with your provider (age 18 and older)								
Hepatitis C Infection	Recommended in adults aged 18-79 years, and in all pregnant women, during each pregnancy									
High Blood Pressure	Recommended every year (age 18 and older)									
HIV Infection	Recommended once (age 18 and older); in all pregnant Veterans; annually in Veterans with ongoing risk factors									
Lung Cancer	Not recommended (age 18_70)						ot recommended ge 81 and older)			
Osteoporosis		Recommended for some won	er (age 18–64)	Recommended once (age 65 and older)						
Sexually Transmitted Infections	Recommended. Test for gonorrhea and chlamydia (age 18–24).  Recommended for some women, talk with your provider about testing for gonorrhea, chlamydia, and syphilis (age 25 and older)									
	Talk with your provider about Recommended for all pregnant Veterans, early screening for syphilis infection, regardless of age. syphilis testing.									

### **MEDICATIONS**

HEALTH CONDITIONS	18-29 years	30–39 years	40–49 years	50-59 years	60–69 years	70-79 years	80 years and older	
Folic Acid for Pregnancy Planning	Recommended. D	Paily folic acid supplement fo pregnant (age 18		Not recommended after child-bearing age (age 55 and older)				
Aspirin to Prevent Cardiovascular Disease	Not recommended (age 18–39)  Talk with your provider (age 40–59)				Not recommended (age 60 and older)			
PrEP to Prevent HIV Infection	Talk with your provider (age 19 and older)							
Statin to Prevent Cardiovascular Disease	Not recommended (age 19–40)  Recommended for some women — talk with your provider (age 40 and older)							

# **HEALTH COUNSELING**

HEALTH CONDITIONS	18–29 years	30–39 years	40–49 years	50–59 years	60–69 years	70–79 years	80 years and older		
Tobacco Use		Recommended every visit (if using tobacco) (age 18 and older)							
Alcohol Use	Talk with your provider about healthy alcohol use (age 18 and older)								
Healthy Diet and Physical Activity		Talk with your provider about a healthy diet and physical activity (age 18 and older)							
Overweight & Obesity	Recommended every year (age 18 and older)								

					// \ 3	<b>,</b>					
VACCINES											
HEALTH CONDITIONS	19–29 years	30–39 years	40–49 yea	ars	50–59 years	60–69 years	70–79 ye	ars	80 years and older		
COVID-19		Recommended for all women — talk with your provider to make sure you stay up-to-date (https://www.va.gov/health-care/covid-19-vaccine/)									
Hepatitis A		Recommended for some women — talk with your provider (age 19 and older)									
Hepatitis B	Recommend	Recommended for all women who have not already had Hepatitis B vaccine (age 19-59)  Recommended for some women - talk with your provider (age 60 and older)									
Herpes Zoster (Shingles) RZV vaccine	Recommended for so	Recommended for some women — talk with your provider (age 19 and older)  Recommended 2 doses (age 50 and older), with second dose 2—6 months af						ths after first dose			
Human Papilloma Virus (HPV) HPV9 vaccine	2–3 doses (age 19–26) if series not completed	Not recommended (age	Not recommended (age 46 and older)								
Influenza (Flu) Injectable, Inhaled				Recomm	nended every year (age 19 a	and older)					
Measles, Mumps, and Rubella (MMR)		Recommended for some women — talk with your provider (age 19 and older)									
Meningococcal Disease		Recommended for some women — talk with your provider (age 19 and older)									
Pneumococcal Disease PCV20 vaccine, PCV15 vaccine, PPSV23 vaccine	Recommended for some women — talk with your provider (age 19—64)  Recommended — talk with your provider to make sure you are up-to-date (age 65 and older)										
<b>Respiratory Syncytial Virus (RSV)</b> RSV vaccine	Pregnant Veterans are recommended to receive RSV Abrysvo® solution between 32-36 weeks gestation during RSV season				vaccine	Recommended for some women — talk with your provider (age 60–74)			mmended for women age 75 and older		
Tdap (Tetanus, Diphtheria & Pertussis)	Recommend at least 1 dose. Get additional dose with every pregnancy (Or for wound management if $>$ 5 years have passed since last tetanus toxoid-containing booster vaccine).										
Td (Tetanus and diphtheria)		Recommended. Booster every 10 years (age 19 and older) (Either Tdap or Td may be used) (Or for wound management if $>$ 5 years have passed since last tetanus toxoid-containing booster vaccine).									
Varicella (Chickenpox)		ed for some women - talk with (2 doses if born 1980 or later)									



provider (2 doses if born 1980 or later)